

Double Transverse-Spin Asymmetries for Small- Q_T Drell-Yan Pair Production in $p\bar{p}$ Collisions

Hiroyuki Kawamura* and Kazuhiro Tanaka†

*Department of Mathematical Sciences, University of Liverpool, Liverpool L69 3BX, UK

†Department of Physics, Juntendo University, Inba, Chiba 270-1695, Japan

Abstract. We discuss the double-spin asymmetries in transversely polarized Drell-Yan process, calculating all-order gluon resummation corrections up to the next-to-leading logarithmic accuracy. This resummation is relevant when the transverse-momentum Q_T of the produced lepton pair is small, and reproduces the (fixed-order) next-to-leading QCD corrections upon integrating over Q_T . The resummation corrections in $p\bar{p}$ -collision behave differently compared with pp -collision cases, and are small at the kinematics in the proposed GSI experiments. This fact allows us to predict large value of the double-spin asymmetries at GSI, using recent empirical information on the transversity.

Keywords: Transversity, Drell-Yan process, Antiprotons, Soft gluon resummation

PACS: 12.38.-t, 12.38.Cy, 13.85.Qk, 13.88.+e

The double-spin asymmetry in Drell-Yan (DY) process with transversely-polarized protons and antiprotons, $p^\uparrow \bar{p}^\uparrow \rightarrow l^+ l^- X$, for azimuthal angle ϕ of a lepton measured in the rest frame of the dilepton $l^+ l^-$ with invariant mass Q and rapidity y , is given by

$$A_{TT} = \frac{d\sigma^{\uparrow\uparrow}/d\omega - d\sigma^{\uparrow\downarrow}/d\omega}{d\sigma^{\uparrow\uparrow}/d\omega + d\sigma^{\uparrow\downarrow}/d\omega} \equiv \frac{\Delta_T d\sigma/d\omega}{d\sigma/d\omega} = \frac{\cos(2\phi)}{2} \frac{\sum_q e_q^2 \delta q(x_1, Q^2) \delta q(x_2, Q^2) + \cdots}{\sum_q e_q^2 q(x_1, Q^2) q(x_2, Q^2) + \cdots}, \quad (1)$$

where $d\omega \equiv dQ^2 dy d\phi$, the summation is over all quark and antiquark flavors with $\delta q(x, Q^2)$ and $q(x, Q^2)$ being the transversity and unpolarized quark-distributions inside a proton, and the ellipses stand for the corrections of next-to-leading order (NLO) and higher in QCD perturbation theory. The scaling variables $x_{1,2}$ represent the momentum fractions associated with the partons annihilating via the DY mechanism, such that $Q^2 = (x_1 P_1 + x_2 P_2)^2 = x_1 x_2 S$ and $y = (1/2) \ln(x_1/x_2)$, where $S = (P_1 + P_2)^2$ is the CM energy squared of $p^\uparrow \bar{p}^\uparrow$. In the proposed polarization experiments at GSI [1], moderate energies, $30 \lesssim S \lesssim 200 \text{ GeV}^2$, allow us to measure (1) for $0.2 \lesssim Q/\sqrt{S} \lesssim 0.7$, and probe the product of the two quark-transversities in the “valence region”. Recently, QCD corrections in (1) at GSI kinematics have been studied: the NLO ($O(\alpha_s)$) corrections [2] as well as the higher order ones in the framework of threshold resummation [3] are rather small, so that the LO value of A_{TT} , which turns out to be large, is rather robust.

When the transverse momentum Q_T of the final $l^+ l^-$ is also observed, we obtain the new double-spin asymmetry as the ratio of the Q_T -differential cross sections, $\mathcal{A}_{TT}(Q_T) \equiv [\Delta_T d\sigma/d\omega dQ_T]/[d\sigma/d\omega dQ_T]$. The bulk of $l^+ l^-$ pair is produced at small $Q_T \ll Q$, where the cross sections $(\Delta_T) d\sigma/d\omega dQ_T$ receive large perturbative corrections with logarithms $\ln(Q^2/Q_T^2)$ multiplying α_s at each order, by the recoil from gluon

radiations, and those have to be treated in an all-order resummation in QCD [4]. The corresponding “ Q_T -resummation” has formally some resemblance to the threshold resummation [3], but embodies contributions from different “edge region” of phase space. The Q_T -resummation for $\Delta_T d\sigma/d\omega dQ_T$ has been derived recently [5], summing the corresponding large logarithms up to next-to-leading logarithmic (NLL) accuracy. Combined with that for $d\sigma/d\omega dQ_T$ [4], we get [6] ($b_0 \equiv 2e^{-\gamma_E}$ with γ_E the Euler constant)

$$\mathcal{A}_{TT}(Q_T) = \frac{\cos(2\phi)}{2} \frac{\int d^2b e^{i\mathbf{b}\cdot\mathbf{Q}_T} e^{S(b,Q)} \sum_q e_q^2 \delta q(x_1, b_0^2/b^2) \delta q(x_2, b_0^2/b^2) + \dots}{\int d^2b e^{i\mathbf{b}\cdot\mathbf{Q}_T} e^{S(b,Q)} \sum_q e_q^2 q(x_1, b_0^2/b^2) q(x_2, b_0^2/b^2) + \dots}, \quad (2)$$

where the numerator and denominator are, respectively, reorganized in the impact parameter b space in terms of the Sudakov factor $e^{S(b,Q)}$ resumming soft and flavor-conserving collinear radiation, while the ellipses involve the remaining contributions of the $O(\alpha_s)$ collinear radiation, which can be absorbed into the exhibited terms as $\delta q \rightarrow \Delta_T C_{qq} \otimes \delta q$, $q \rightarrow C_{qq} \otimes q + C_{qg} \otimes g$ using the corresponding coefficient functions $(\Delta_T)C_{ij}$; note that there is no “chiral-odd” gluon distribution to participate in the numerator of (2) as well as (1). Using *universal* Sudakov exponent $S(b, Q)$ with the first nonleading anomalous dimensions in (2), the first three towers of large logarithmic contributions to the cross sections, $\alpha_s^n \ln^m(Q^2/Q_T^2)/Q_T^2$ ($m = 2n - 1, 2n - 2, 2n - 3$), are resummed to all orders in α_s , yielding the NLL resummation. In addition to these resummed components relevant for small Q_T , the ellipses in (2) also involve the other terms of the fixed-order α_s , which treat the LO processes in the large Q_T region, so that (2) is the ratio of the NLL+LO polarized and unpolarized cross sections. We include a Gaussian smearing as usually as $S(b, Q) \rightarrow S(b, Q) - g_{NP} b^2$, corresponding to intrinsic transverse momentum of partons inside proton. The integrations of the NLL+LO cross sections $\Delta_T d\sigma/d\omega dQ_T$ and $d\sigma/d\omega dQ_T$ over Q_T coincide, respectively, with the (fixed-order) NLO cross sections $\Delta_T d\sigma/d\omega$ and $d\sigma/d\omega$, associated with A_{TT} of (1) [5, 6].

The resummation makes $1/b \sim Q_T$ the relevant scale in (2), in contrast to Q in (1). Figure 1 shows [6] the numerical evaluation of (2), as well as of $\Delta_T d\sigma/d\omega dQ_T$ associated with its numerator, at GSI kinematics using the NLO transversities that saturate the Soffer bound, $2\delta q(x, \mu^2) \leq q(x, \mu^2) + \Delta q(x, \mu^2)$, at a low scale μ with Δq the helicity distribution. The NLL resummed component dominates $\Delta_T d\sigma/d\omega dQ_T$ in small and moderate Q_T region, and similarly for $d\sigma/d\omega dQ_T$ reflecting universality of the large Sudakov effects, which leads to almost constant $\mathcal{A}_{TT}(Q_T)$, in particular, with even flatter behavior than the corresponding asymmetry [5] for the pp -collision case. Remarkably, $\mathcal{A}_{TT}(Q_T)$ at NLL+LO has almost the same value as that at LL; this is in contrast to the pp case where the resummation at higher level enhances the asymmetry [5]. We note that $\mathcal{A}_{TT}(Q_T)$ at LL is given by (2) omitting all nonleading corrections, i.e., omitting the ellipses, replacing $S(b, Q)$ by that at the LL level, and replacing the scale of the parton distributions as $b_0^2/b^2 \rightarrow Q^2$, so that the result coincides with A_{TT} at LO (see (1)). Therefore, at GSI, both $\mathcal{A}_{TT}(Q_T)$ and A_{TT} are quite stable when including the QCD (resummation and fixed-order) corrections, with $\mathcal{A}_{TT}(Q_T) \simeq \mathcal{A}_{TT}(0)$, and

$$\mathcal{A}_{TT}(Q_T) \simeq A_{TT}. \quad (3)$$

To clarify the relevant mechanism, $\mathcal{A}_{TT}(Q_T) \simeq \mathcal{A}_{TT}(0)$ allows us to consider the $Q_T \rightarrow 0$ limit of (2): for $Q_T \approx 0$, the b integral is controlled by a saddle point $b = b_{SP}$,




FIGURE 1. (a) $\Delta_T d\sigma/dQ^2 dy dQ_T$ (b) $\mathcal{A}_{TT}(Q_T)$ with GSI kinematics, $S = 210 \text{ GeV}^2$, $Q = 4 \text{ GeV}$, $y = \phi = 0$, and with $g_{NP} = 0.5 \text{ GeV}^2$, using the NLO transversity distributions that correspond to the Soffer bound.

which has the same value between the numerator and denominator in (2) such that [5, 6]

$$\mathcal{A}_{TT}(0) \simeq \frac{\cos(2\phi)}{2} \frac{\sum_q e_q^2 \delta q(x_1, b_0^2/b_{SP}^2) \delta q(x_2, b_0^2/b_{SP}^2)}{\sum_q e_q^2 q(x_1, b_0^2/b_{SP}^2) q(x_2, b_0^2/b_{SP}^2)}, \quad (4)$$

omitting the small corrections from the LO components involved in the ellipses in (2). This saddle-point evaluation is exact at NLL accuracy; in particular, the $O(\alpha_s)$ contributions from the coefficients $(\Delta_T)C_{ij}$, e.g., $C_{qg} \otimes g$ associated with gluon distribution, completely decouple as $Q_T \rightarrow 0$ (see [4, 5]). The simple form of (4) is reminiscent of A_{TT} of (1) at LO, but is different from the latter in the unconventional scale b_0^2/b_{SP}^2 ; the actual position of the saddle point implies $b_0/b_{SP} \simeq 1 \text{ GeV}$, irrespective of the values of Q and g_{NP} [5, 6]. In the valence region relevant for GSI kinematics, the u -quark contribution dominates in (4) and (1), so that these asymmetries are controlled by the ratio, $\delta u(x_{1,2}, \mu^2)/u(x_{1,2}, \mu^2)$ with $\mu^2 = b_0^2/b_{SP}^2$ and Q^2 , respectively. Actually the scale dependence in this ratio almost cancels between the numerator and denominator as $\delta u(x, b_0^2/b_{SP}^2)/u(x, b_0^2/b_{SP}^2) \simeq \delta u(x, Q^2)/u(x, Q^2)$ (see Fig. 3 in [6]); this implies (3) at GSI. Note that this is not the case for pp collisions because of very different behavior of the sea-quark components under the evolution between transversity and unpolarized distributions [5]; indeed $\mathcal{A}_{TT}(Q_T) > A_{TT}$ at RHIC and J-PARC [5]. A similar logic applied to (2) also explains why $\mathcal{A}_{TT}(Q_T)$ at GSI are flatter than in pp collisions.

Another consequence of the similar logic is that $\delta u(x_{1,2}, 1 \text{ GeV}^2)/u(x_{1,2}, 1 \text{ GeV}^2)$ as a function of $x_{1,2}$ directly determines the Q - as well as S -dependence of the value of (3) at GSI, through $x_{1,2} = (Q/\sqrt{S})e^{\pm y}$. In Fig. 2, using the NLO transversity distributions corresponding to the Soffer bound, the symbols “ Δ ” plot $\mathcal{A}_{TT}(Q_T \simeq 1 \text{ GeV})$ of (2) at NLL+LO [6]. The dashed curve draws the result using (4); this simple formula indeed works well. Also plotted by the two-dot-dashed curve is A_{TT} of (1) at LO with the transversities corresponding to the Soffer bound at LO level, to demonstrate (3). The Q - and S -dependence of these results reflect that the ratio $\delta u(x, 1 \text{ GeV}^2)/u(x, 1 \text{ GeV}^2)$ is an increasing function of x . These results using the Soffer bound show the “maximally possible” asymmetry, i.e., optimistic estimate. A more realistic estimate of (2) (with $Q_T \simeq 1 \text{ GeV}$) and (4) is shown [6] in Fig. 2 by the symbols “ ∇ ” and the dot-dashed curve, respectively, with the NLO transversity distributions assuming $\delta q(x, \mu^2) = \Delta q(x, \mu^2)$ at




FIGURE 2. The double transverse-spin asymmetries at GSI as functions of Q with $y = \phi = 0$.

a low scale μ , as suggested by nucleon models and favored by the results of empirical fit for transversity [7]. The new estimate gives smaller asymmetries compared with the Soffer bound results, but still yields rather large asymmetries [6]. Based on (3), these results in Fig. 2 may be considered as estimate of A_{TT} of (1).

At present, empirical information of transversity is based on the LO global fit, using the semi-inclusive DIS data and assuming that the antiquark transversities vanish, $\delta\bar{q}(x) = 0$, so that the corresponding LO parameterization is available only for u and d quarks [7]. Fortunately, however, the dominance of the u -quark contribution in the GSI kinematics allows quantitative evaluation of A_{TT} at LO using only this empirical information [6]: the upper limit of the error band for the u - and d -quark transversities obtained by the global fit [7] yields the “upper bound” of A_{TT} shown by the dotted curve in Fig. 2; using (3), this result may be considered also as estimate of $\mathcal{A}_{TT}(Q_T)$. In the small Q region, our full NLL+LO result of $\mathcal{A}_{TT}(Q_T \simeq 1 \text{ GeV})$, shown by “ ∇ ”, can be consistent with estimate using the empirical LO transversity, but these results have rather different behavior for increasing Q , because the u -quark transversity for the former lies slightly outside the error band of the global fit for $x \gtrsim 0.3$ [6]. Thus, in the large asymmetries to be observed at GSI, the behavior of $\mathcal{A}_{TT}(Q_T)$, A_{TT} as functions of Q will allow us to determine the detailed shape of transversity distributions.

ACKNOWLEDGMENTS

This work was supported by the Grant-in-Aid for Scientific Research No. B-19340063.

REFERENCES

1. V. Barone *et al.* [PAX Collaboration], arXiv:hep-ex/0505054.
2. V. Barone, A. Caferella, C. Coriano, M. Guzzi and P. G. Ratcliffe, Phys. Lett. **B639**, 483 (2006).
3. H. Shimizu, G. Sterman, W. Vogelsang and H. Yokoya, Phys. Rev. **D71**, 114007 (2005).
4. J. C. Collins, D. E. Soper and G. Sterman, Nucl. Phys. **B250**, 199 (1985).
5. H. Kawamura, J. Kodaira, H. Shimizu and K. Tanaka, Prog. Theor. Phys. **115**, 667 (2006); H. Kawamura, J. Kodaira and K. Tanaka, Nucl. Phys. **B777**, 203 (2007); H. Kawamura, J. Kodaira and K. Tanaka, Prog. Theor. Phys. **118**, 581 (2007).
6. H. Kawamura, J. Kodaira and K. Tanaka, Phys. Lett. **B662**, 139 (2008).
7. M. Anselmino *et al.*, Phys. Rev. **D75**, 054032 (2007); arXiv:0812.4366 [hep-ph].